



TITLE:

Microlocal analysis on PDE : Some contributions by Yoshinori Morimoto around kinetic equations (Workshop on the Boltzmann Equation, Microlocal Analysis and Related Topics)

AUTHOR(S):

Xu, Chao-Jiang

---

CITATION:

Xu, Chao-Jiang. Microlocal analysis on PDE : Some contributions by Yoshinori Morimoto around kinetic equations (Workshop on the Boltzmann Equation, Microlocal Analysis and Related Topics). 数理解析研究所講究録別冊 2017, B67: 137-154

ISSUE DATE:

2017-10

URL:

<http://hdl.handle.net/2433/243709>

RIGHT:

© 2017 by the Research Institute for Mathematical Sciences, Kyoto University. All rights reserved.

# Microlocal analysis on PDE : Some contributions by Yoshinori Morimoto around kinetic equations

By

Chao-Jiang Xu \*

## Abstract

In this short paper, we give a brief presentation of some contribution to microlocal analysis on the partial differential equations by Yoshinori Morimoto. The first part concerns the study of the degenerate elliptic equations, by using the microlocal analysis based on the theory of pseudo-differential operators. The second part is about the analysis of non-cutoff Boltzmann equations where the microlocal analysis contribute lots of progress, in particular, using Fefferman-Phong's uncertainty principle to prove the smoothing effect of solutions and Littlewood-Paley theory to study the existence of classical solutions for non-cutoff Boltzmann equations. So that we focus only on the contribution to the study of kinetic equations by Yoshinori Morimoto.

## § 1. Hypoellipticity of infinite degenerate elliptic operators

*Microlocal analysis (Wikipedia Encyclopedia)*

*In mathematical analysis, microlocal analysis comprises techniques developed from the 1950s onwards based on Fourier transforms related to the study of linear PDE. This includes pseudo-differential operators and Fourier integral operators. The term microlocal implies localisation not only with respect to location in the space, but also with respect to cotangent space directions at a given point. The techniques of microlocal*

---

Received December 22, 2016. Revised May 23, 2017.

2010 Mathematics Subject Classification(s): 35A05, 35B65, 35D10, 35H20, 76P05, 84C40

*Key Words:* Microlocal analysis, hypoellipticity, uncertainty principle, Littlewood-Paley theory, non-cutoff Boltzmann equations.

\*Université de Rouen, CNRS, UMR 6085-Laboratoire de Mathématiques 76801 Saint-Etienne du Rouvray, France

and

School of Mathematics and Statistics, Wuhan University 430072, Wuhan, P. R. China

e-mail: [chao-jiang.xu@univ-rouen.fr](mailto:chao-jiang.xu@univ-rouen.fr)

© 2017 Research Institute for Mathematical Sciences, Kyoto University. All rights reserved.

*analysis were developed in the 1960s and 1970s.*

**Hypoellipticity :** A partial differential operator  $P$  is called hypoelliptic, if  $u \in \mathcal{D}', Pu \in C^\infty$  imply that  $u \in C^\infty$ .

- Elliptic operator with  $C^\infty$  coefficients is hypoelliptic.
- The following Hörmander's operators is hypoelliptic,

$$H = \sum_{j=1}^m X_j^2 + X_0,$$

where  $X = \{X_0, X_1, \dots, X_m\}$  is a system of vector fields which satisfy the Hörmander's condition:  $X_0, X_1, \dots, X_m$  and their repeated commutators of at most  $k$  times span the tangent space .

A typic exemple of Hörmander's operators is

$$P_1 = \partial_t^2 + \partial_x^2 + x^{2k} \partial_y^2.$$

It is possible to take a system of pseudo-differential operators  $X$ .

The work of Y. Morimoto was motivated by the previous result of S. Kusuoka and D. Stroock by means of the probabilistic method ( Malliavin calculus). Namely, they gave a remarkable example:

$$P = D_t^2 + D_x^2 + g(x)D_y^2, \quad 0 \leq g(x) \in C^\infty(\mathbb{R}), \quad g(x) > 0 \text{ if } x \neq 0$$

is hypoelliptic in  $\mathbb{R}^3$ , if and only if

$$\lim_{x \rightarrow \infty} |x| \log g(x) = 0.$$

If  $g(x) = \exp -|x|^\sigma, \sigma > 0$  then the condition is interpreted to  $0 < \sigma < 1$ . The term  $D_t^2$  is significant, in fact,  $A = D_x^2 + g(x)D_y^2$  is hypoelliptic in  $\mathbb{R}^2$  without the degenerate rate condition. This phenomena is quite similar to non-analytic hypoellipticity for Baouendi-Goulaouic operator

$$P_0 = D_t^2 + D_x^2 + x^{2m} D_y^2, \quad m = 1, 2, \dots,$$

that is, the Gevrey singularity of order  $G^{m+1}$  propagates along the  $t$  direction. The propagation of the singularity of Gevrey class can be roughly understood by the uncertainty principle ( mathematically formalized by Fefferman-Phong) as follows: If we denote the symbol of  $D_x^2 + x^{2m} D_y^2$  by  $\xi^2 + x^{2m} \eta^2$ , then

$$0 \leq \xi^2 + x^{2m} \eta^2 \leq 2|\eta|^{2/(m+1)} \text{ on the box } \{|x| \leq |\eta|^{-1/(m+1)}, |\xi| \leq |\eta|^{1/(m+1)}\},$$

and  $P_0$  behaves like, on a tube  $\{(t, x, y, \xi, \eta); t \in (-\infty, \infty), |x| \leq |\eta|^{-1/(m+1)}, |\xi| \leq |\eta|^{1/(m+1)}\}$ ,

$$P_0 \sim D_t^2 + 2|D_y|^{2/(m+1)}.$$

As for  $C^\infty$  singularity, the log order is critical. Based on this idea, the characterization of partial differential operators with the logarithmic regularity estimate has been studied by means of the Fefferman-Phong uncertainty principle.

**Infinite degenerate elliptic operators** The following example is an infinite degenerate elliptic operators

$$P_2 = \partial_t^2 + \partial_x^2 + e^{-2|x|^{-\delta}} \partial_y^2,$$

with  $0 < \delta < 1$ . This operator is also hypoelliptic. More generally, for the operators

$$P_3 = D_t^2 + \mathcal{A}(x, D_x), \quad \mathbb{R}_t \times \mathbb{R}_x^n$$

where  $\mathcal{A}(x, D_x)$  is a formally self-adjoint second order PDO such that

$$(\mathcal{A}(x, D_x)u, u) \geq c_0 \|u\|_{L^2}^2, \quad \forall u \in C_0^\infty.$$

There are a criterion for hypoellipticity.

*Theorem 1.1* (Y. Morimoto, Osaka J. Math. 1987 ).  $P_3$  is hypoelliptic in  $\mathbb{R}_t \times \mathbb{R}_x^n$  if and only if for any  $x_0 \in \mathbb{R}_x^n$  there exists a neighborhood  $\omega$  of  $x_0$  such that for any  $\epsilon > 0$  the estimate

$$\|\log \langle D_t, D_x \rangle \phi\|_{L^2}^2 \leq \epsilon \operatorname{Re}(P_3 \phi, \phi) + C_\epsilon \|\phi\|_{L^2}^2, \quad \forall \phi \in C_0^\infty(\mathbb{R}_t \times \omega)$$

holds with a constant  $C_\epsilon$ .

The above results can be extended to more general cases.

**Another necessary and sufficient condition for the hypoellipticity** Let

$$P_4 = D_x^2 + f(x)D_t^2 + g(x)D_y^2, \quad \mathbb{R}_{x,t,y}^3, \quad f, g \geq 0.$$

Let  $f_I$  denote the average of  $f$  in  $I \subset I_0$  an open fixed interval in  $\mathbb{R}_x$ ,  $f_I, g_I > 0$ . We say that  $f, g$  satisfy  $(M, f, g)$  condition in  $I_0$  if

$$(M, f, g) \quad \inf_{\delta > 0} \left( \sup \{ f_I^{1/2} |I| |\log g_{3I}|; I \text{ with } 3I \subset I_0, g_{3I} < \delta \} \right) = 0.$$

*Theorem 1.2* (Y. Morimoto-T. Morioka, Bull. Sci. Math. 1997 ).  $P_4$  is hypoelliptic in  $I_0 \times \mathbb{R}_t \times \mathbb{R}_y$  if and only if  $f, g$  satisfy  $(M, f, g)$  and  $(M, g, f)$  in  $I_0$ .

The above results is also true for more general cases. An example :

$$f = x^{2m}, \quad g = e^{-|x|^{-s}}, \quad s < m + 1, \text{ (by } T.Hoshiro \text{)}$$

**The uncertainty principle** Consider a symbol of the form, for  $0 < \lambda \leq 1$ ,

$$a(x, \xi) = |\xi|^{2\lambda} + V(x), \quad (x, \xi) \in \mathbb{R}^{2n}$$

where  $V(x)$  is a non-negative function and depends on a large parameter  $M > 0$ . Let  $\mathcal{C}$  denote the set of boxes

$$B = \{(x, \xi) \in \mathbb{R}^{2n}; |x_j - x_j^0| \leq \frac{\delta}{2}, |\xi_j - \xi_j^0| \leq \frac{\delta^{-1}}{2}\}.$$

*Theorem 1.3* (Y. Morimoto, Publ. RIMS. Kyoto Univ. 1987).

$$\inf_{B \in \mathcal{C}} m(\{(x, \xi) \in B; a(x, \xi) \geq R\}) \geq c > 0,$$

imply

$$\| |D_x|^\lambda u \|_{L^2}^2 + (V(x)u, u) \geq c'R \|u\|_{L^2}^2, \quad u \in C_0^\infty(\mathbb{R}^n),$$

where  $c' > 0$  depends only on  $c, n, \lambda$ .

As an application, we consider the operators

$$P_5 = a(x, y, D_x) + g(x')b(x, y, D_y), \quad \text{in } \mathbb{R}^n \times \mathbb{R}^m,$$

where  $x = (x', x'') \in \mathbb{R}^n$  and

$$\begin{aligned} \operatorname{Re} a(x, y, \xi) &\geq c_1 |\xi|^2, \quad \text{for large } |\xi|, \\ \operatorname{Re} b(x, y, \xi) &\geq c_2 |\eta|^2, \quad \text{for large } |\eta|, \end{aligned}$$

$g(x') > 0$  for all  $x' \neq 0$  and vanish infinite order at 0. Then we have

$$\lim_{|x'| \rightarrow 0} |x'| |\log g(x')| = 0 \implies P_5 \text{ is hypoelliptic.}$$

Let  $a(x, \xi) = |\xi|^2 + c g(x') |\eta|^2$ , then we can apply the uncertainty principle with  $R = (k \log |\eta|)^2$  which deduce the logarithmic regularity estimates.

## § 2. The non-cutoff Boltzmann equations

From 2005, Yoshinori Morimoto and his collaborators (Alexandre-Morimoto-Ukai-Xu-Yang (AMUXY)) start to work on the Kinetic equations such as non-cutoff Boltzmann equations and Landau equation with more than 30 publications.

$$(2.1) \quad f_t + v \cdot \nabla_x f = Q(f, f), \quad f|_{t=0} = f_0,$$

where the Boltzmann collisional operator is

$$Q(g, f) = \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} B(v - v_*, \sigma) \{g'_* f' - g_* f\} d\sigma dv_*,$$

and the collision kernel is

$$B(v - v_*, \sigma) = |v - v_*|^\gamma b(\cos \theta), \quad 0 \leq \theta \leq \frac{\pi}{2},$$

with angular non-cutoff factor,

$$(2.2) \quad b(\cos \theta) \approx \theta^{-2-2s}, \quad 0 \leq \theta \leq \frac{\pi}{2}.$$

### Smoothing effect of non-cutoff Boltzmann equation

A first microlocal analysis results of Boltzmann equation is the following smoothing effect work.

*Theorem 2.1* (AMUXY, ARMA, 2010). Assume (2.2) holds true, with  $0 < s < 1$ ,  $\gamma > \max\{-3, -3/2 - 2s\}$ ,  $0 < T \leq +\infty$ .  $\Omega \subset \mathbb{R}_x^3$ ,  $f \geq 0$ ,  $f \not\equiv 0$ ,  $f \in L^\infty([0, T]; H_\ell^5(\Omega \times \mathbb{R}^3))$  for any  $\ell \in \mathbb{N}$ , be a solution of Cauchy problem (2.1). Then we have

$$f \in C^\infty([0, T] \times \Omega, \mathcal{S}(\mathbb{R}^3)).$$

The main ingredient for the proof of the above Theorem are :

- The coercivity and upper bound estimates of Boltzmann operators.
- Uncertainty principle and regularity of the following transport equation:

$$(2.3) \quad f_t + v \cdot \nabla_x f = g \in \mathcal{D}'(\mathbb{R}^7).$$

**The coercivity and upper bound estimates** We have the following coercive estimate :

Let  $g \geq 0$ ,  $g \not\equiv 0$ ,  $g \in L_2^1 \cap L \log L(\mathbb{R}_v^3)$ . Then for any function  $f \in H_{\frac{\gamma}{2}+s}^s(\mathbb{R}_v^3)$ , we have

$$C_g^{-1} \|f\|_{H_{\frac{\gamma}{2}}^s(\mathbb{R}_v^3)}^2 \leq (-Q(g, f), f)_{L^2(\mathbb{R}_v^3)} + C \|g\|_{L_{\gamma+2s}^1(\mathbb{R}_v^3)} \|f\|_{L_{\frac{\gamma}{2}+s}^2(\mathbb{R}_v^3)}^2.$$

We have also the upper bound estimate

$$\|Q(f, g)\|_{H_\alpha^m(\mathbb{R}_v^3)} \leq C \|f\|_{L_{\alpha+(\gamma+2s)+}^1(\mathbb{R}_v^3)} \|g\|_{H_{(\alpha+\gamma+2s)+}^{m+2s}(\mathbb{R}_v^3)}.$$

The results are due to : Alexandre-Desvillettes-Villani-Wennberg and also Alexandre-Morimoto-Ukai-Xu-Yang (AMUXY).

### The Littlewood-Paley theory

- For the coercivity and upper bound estimate of non-cutoff Boltzmann operators, in the Maxwellian case ( $\gamma = 0$ ), the proof is just by using the Fourier transformation (Bobylov formula).
- But for general non Maxwellian case ( $\gamma > -3$ ), we need to use the following Littlewood-Paley decomposition :

$$\phi_0(v) + \sum_{k=1}^{\infty} \phi(2^{-k}v) = 1, \quad v \in \mathbb{R}^3,$$

with  $\phi_0, \phi \in C_0^\infty(\mathbb{R}^3)$  and

$$\text{supp } \phi_0 \subset \{|v| < 2\}, \quad \text{supp } \phi \subset \{1 < |v| < 3\}.$$

Then we can study the Boltzmann operators with kinetic factors as

$$B_k(|v - v_*|, \sigma) = \phi(2^{-k}(v - v_*))|v - v_*|^\gamma b(\cos \theta).$$

### The Fefferman-Phong's uncertainty principle and kinetic equations

The smoothing property of Boltzmann equation is like the following Hörmander's type equation:

$$f_t + v \cdot \nabla_x f + (-\Delta_v)^s f = g,$$

with  $X_0 = \partial_t + v \cdot \nabla_x$ ,  $X_1 = (-\Delta_v)^{\frac{s}{2}}$ . And it is in the form of transport equation as (2.3).

Let  $a(r) = r^{-2s}$ ,  $Q$  is a cube in  $\mathbb{R}^n$ ,  $\ell(Q)$  is the side length.  $Q^*$  is any cube such that  $Q \subset Q^*$  with  $\ell(Q^*) = 2\ell(Q)$ . For  $a^+, a^- \geq 0$ ,

$$E(Q, Q^*) = \{v \in Q^*; a^+(v) \geq \|a^-\|_{L^\infty(Q)} - a(\ell(Q))\}.$$

The main assumption on the functions  $a^+, a^-$  is

$$(H_0) \quad \inf_{Q, Q^*} \left\{ \frac{|E(Q, Q^*)|}{|Q^*|}; \quad Q \subset Q^* \text{ with } 2\ell(Q) = \ell(Q^*) \right\} \geq \kappa > 0.$$

### Regularity for transport Equations

*Theorem 2.2* ( Uncertainty Principle, JFA, AMUXY, 2008). If the condition  $(H_0)$  holds, then

$$\int_{\mathbb{R}^n} a^-(v) |f(v)|^2 dv \leq C \int_{\mathbb{R}^n} (|D_v|^s f(v)|^2 + a^+(v) |f(v)|^2) dv,$$

for any  $f \in \mathcal{S}(\mathbb{R}^n)$ .

*Theorem 2.3* (JFA, AMUXY, 2008). Assume that  $g \in H^{-s'}(\mathbb{R}^7)$ ,  $0 \leq s' < 1$ ,  $f \in L^2(\mathbb{R}^7)$  is a weak solution (2.3) such that  $|D_v|^s f \in L^2(\mathbb{R}^7)$  for some  $0 < s \leq 1$ . Then

$$\frac{\Lambda_x^{s(1-s')/(s+1)} f}{(1 + |v|^2)^{s s'/2(s+1)}}, \quad \frac{\Lambda_t^{s(1-s')/(s+1)} f}{(1 + |v|^2)^{s/2(s+1)}} \in L^2(\mathbb{R}^7).$$

The main idea in the proof of this regularity theorem is to define the corresponding functions  $a_{\pm}(v)$  and show that they satisfy the condition  $(H_0)$  with the Fourier variables  $\tau$  and  $\eta$  as parameters.

$$a^-(v, \tau, \eta) = c_0 \frac{|\tau|^{2s(1-s')/(s+1)}}{(1 + |v|^2)^{s/(s+1)}} + c_1 \frac{|\eta|^{2s(1-s')/(s+1)}}{(1 + |v|^2)^{s s'/2(s+1)}},$$

and

$$a^+(v, \tau, \eta) = 1 + \frac{|\tau + v \cdot \eta|^2}{(1 + \tau^2 + |\eta|^2)^{s'}}.$$

### Existence of the classical solutions

Consider the perturbation  $f = \mu + \sqrt{\mu}g$  around a normalized Maxwellian distribution

$$\mu(v) = (2\pi)^{-\frac{3}{2}} e^{-\frac{|v|^2}{2}}.$$

Since  $Q(\mu, \mu) = 0$ , we have

$$Q(f, f) = Q(\mu, \sqrt{\mu}g) + Q(\sqrt{\mu}g, \mu) + Q(\sqrt{\mu}g, \sqrt{\mu}g).$$

Denote

$$\Gamma(g, h) = \mu^{-1/2} Q(\sqrt{\mu}g, \sqrt{\mu}h).$$

Then the linearized Boltzmann operator takes the form

$$\mathcal{L}g = -\Gamma(\sqrt{\mu}, g) - \Gamma(g, \sqrt{\mu}).$$

We consider the Cauchy problem for the perturbation  $g$

$$(2.4) \quad \begin{cases} g_t + v \cdot \nabla_x g + \mathcal{L}g = \Gamma(g, g), & t > 0, \\ g|_{t=0} = g_0. \end{cases}$$

We have the following existence of classical solutions for non-cutoff Boltzmann equation.

*Theorem 2.4* (JFA, AMUXY, 2012). Assume  $\gamma + 2s \leq 0$ ,  $0 < s < 1$  and  $\gamma > \max\{-3, -\frac{3}{2} - 2s\}$ . Let  $N \geq 4, \ell \geq N$ . There exists  $\varepsilon_0 > 0$ , such that if  $\|g_0\|_{\mathcal{H}_\ell^N(\mathbb{R}^6)} \leq \varepsilon_0$ , then the Cauchy problem (2.4) admits a global solution

$$g \in L^\infty([0, +\infty[; \mathcal{H}_\ell^N(\mathbb{R}^6)).$$



Here

$$\|f\|_{\mathcal{H}_\ell^N(\mathbb{R}^6)}^2 = \sum_{|\alpha|+|\beta|\leq N} \|\langle v \rangle^{2s+\gamma|\ell-|\beta||} \partial_\beta^\alpha f\|_{L^2(\mathbb{R}^6)}^2.$$

*Theorem 2.5* (AA, AMUXY, 2011). Assume  $0 < s < 1$  and  $\gamma + 2s > 0$ . Let  $g_0 \in H_\ell^k(\mathbb{R}^6)$  for some  $k \geq 6$ ,  $\ell > 3/2 + 2s + \gamma$ . There exists  $\varepsilon_0 > 0$ , such that if  $\|g_0\|_{H_\ell^k(\mathbb{R}^6)} \leq \varepsilon_0$ , the Cauchy problem (2.4) admits a global solution

$$g \in L^\infty([0, +\infty[; H_\ell^k(\mathbb{R}^6)).$$

By apply Theorem 2.1, then we have the regularity of solutions of Theorem 2.4-2.5,

$$f = \mu + \sqrt{\mu} g \in C^\infty([0, T[\times \mathbb{R}^3, \mathcal{S}(\mathbb{R}^3)).$$

### § 3. The smoothing effect of the non-cutoff Boltzmann operators

#### Gelfand-Shilov regularizing effect

Another important work of Y. Morimoto on the microlocal analysis of kinetic equation is the regularizing effect of Cauchy problem in Gelfand-Shilov class.

The Gelfand-Shilov spaces  $S_\nu^\mu(\mathbb{R}^d)$ , with  $\mu, \nu > 0$ ,  $\mu + \nu \geq 1$ , are defined as the spaces of smooth functions  $f \in C^\infty(\mathbb{R}^d)$  satisfying the estimates

$$\exists A, C > 0, \quad |\partial_v^\alpha f(v)| \leq CA^{|\alpha|} (\alpha!)^\mu e^{-\frac{1}{A}|v|^{1/\nu}}, \quad v \in \mathbb{R}^d, \alpha \in \mathbb{N}^d,$$

or, equivalently

$$\exists A, C > 0, \quad \sup_{v \in \mathbb{R}^d} |v^\beta \partial_v^\alpha f(v)| \leq CA^{|\alpha|+|\beta|} (\alpha!)^\mu (\beta!)^\nu, \quad \alpha, \beta \in \mathbb{N}^d.$$

Also equivalently,  $f \in \mathcal{S}(\mathbb{R}^d)$  satisfying to the estimates

$$\exists C > 0, \epsilon > 0, \quad |f(v)| \leq Ce^{-\epsilon|v|^{1/\nu}}, \quad |\widehat{f}(\xi)| \leq Ce^{-\epsilon|\xi|^{1/\mu}}.$$

Remark that Gelfand-Shilov space is much strong than the Gevrey space(for which  $\nu = +\infty$ ) and the Schwartz space  $\mathcal{S}(\mathbb{R}^d)$ .

The Cauchy problem defined by the evolution equation associated to the harmonic oscillator

$$\begin{cases} \partial_t f + \mathcal{H}f = 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

enjoys nice regularizing properties in  $S_{\frac{1}{2}}^{\frac{1}{2}}(\mathbb{R}^d)$  for any positive time, where

$$\mathcal{H} = -\Delta_v + \frac{|v|^2}{4}$$

is the  $d$ -dimensional harmonic oscillator.

Moreover, the smoothing effect for the solutions to the Cauchy problem defined by the evolution equation associated to the fractional harmonic oscillator,  $0 < s < 1$ ,

$$\begin{cases} \partial_t f + \mathcal{H}^s f = 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

occurs for any positive time in the symmetric Gelfand-Shilov space  $S_{\frac{1}{2s}}^{\frac{1}{2s}}(\mathbb{R}^d)$ .

The first result about kinetic equation is the regularizing effect of Cauchy problem to the following Kolmogorov equation :

$$\begin{cases} \partial_t f + v \cdot \nabla_x f - \Delta_v f = 0, & (x, v) \in \mathbb{R}^{2d} \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^{2d}). \end{cases}$$

The solution of above equation satisfying  $f(t) \in G^{\frac{1}{2}}(\mathbb{R}^{2d})$ , for all  $t > 0$ , in fact, we have

$$e^{-c(t\Delta_v + t^3\Delta_x)} f(t, \cdot, \cdot) \in L^2(\mathbb{R}^{2d}),$$

for some  $c > 0$  (Morimoto-Xu [26]).

For the solution of the spatial homogeneous radial symmetric non-cutoff Boltzmann equation (Maxwellian molecule), we consider also a perturbation  $f = \mu + \sqrt{\mu}g$ , then we have

*Theorem 3.1* (Lerner-Morimoto-Pravda-Starov-Xu, JDE, 2013). Let  $g_0 \in L^1(\mathbb{R}_v^3) \cap L^2(\mathbb{R}_v^3)$  radially symmetric and  $\|g_0\|_{L^2} \ll 1$ . Assume that  $g(t)$  is the classical solution of the Cauchy problem

$$\partial_t g + \mathcal{L}g = \Gamma(g, g), \quad g|_{t=0} = g_0.$$

Then,  $\exists c_0 > 0, C > 0, \forall t \geq 0$ ,

$$\|e^{c_0 t \mathcal{H}^s} g(t)\|_{L^2} \leq e^{-Ct} \|g_0\|_{L^2},$$

this implies that

$$\forall t > 0, \quad g(t) \in S_{\frac{1}{2s}}^{\frac{1}{2s}}(\mathbb{R}^3).$$

It is also true for the case  $s = 1$ , i. e. the homogeneous Landau equation.

Main ingredient of proofs of above results is the spectral decomposition of harmonic oscillator on the Hermite basis, where the Hermite function is

$$\Psi_\alpha = \frac{1}{\sqrt{\alpha_1! \dots \alpha_d!}} A_1^{\alpha_1} \dots A_d^{\alpha_d} \Psi_0, \quad \alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}^d,$$

with

$$\Psi_0(v) = (2\pi)^{-\frac{d}{4}} e^{-\frac{|v|^2}{4}}, \quad A_j = \frac{v_j}{2} - \frac{\partial}{\partial v_j}.$$

The family  $(\Psi_\alpha)_{\alpha \in \mathbb{N}^d}$  is an orthonormal basis of  $L^2(\mathbb{R}^d)$  composed by the eigenfunctions of the  $d$ -dimensional harmonic oscillator

$$\mathcal{H} = -\Delta_v + \frac{|v|^2}{4} = \sum_{k \geq 0} \left( \frac{d}{2} + k \right) \mathbb{P}_k, \quad \text{Id} = \sum_{k \geq 0} \mathbb{P}_k,$$

where  $\mathbb{P}_k$  is the orthogonal projection onto  $\mathcal{E}_k = \text{Span}\{\Psi_\alpha; |\alpha| = k\}$ .

For the linearized radially symmetric Boltzmann operators, we have the following spectral decomposition

$$\mathcal{L} = \sum_{k \geq 1} \lambda_k \mathbb{P}_k, \quad \lambda_k \approx k^s, \quad (k \rightarrow +\infty).$$

Then, we have

$$(\mathcal{L}g, g)_{L^2} \approx \|\mathcal{H}^{s/2}g\|_{L^2},$$

and the upper bounded estimate of Boltzmann operators,

$$\left( \Gamma(f, g), e^{t\mathcal{L}} \mathbf{S}_n h \right)_{L^2} \leq C \|e^{\frac{t}{2}\mathcal{L}} \mathbf{S}_{n-2} f\|_{L^2} \|e^{\frac{t}{2}\mathcal{L}} \mathbf{S}_{n-2} \mathcal{H}^{s/2} g\|_{L^2} \|e^{\frac{t}{2}\mathcal{L}} \mathbf{S}_n \mathcal{H}^{s/2} h\|_{L^2}$$

with  $\mathbf{S}_n = \sum_{k=0}^n \mathbb{P}_k$ . This spectral decomposition permits to study also the smoothing effect of the spatially inhomogeneous non-cutoff Kac equation,

$$\begin{cases} \partial_t f + v \partial_x f = K(f, f), \\ f|_{t=0} = f_0, \end{cases}$$

the Kac collision operator is defined as

$$K(g, f) = \int_{|\theta| \leq \frac{\pi}{4}} \beta(\theta) \left( \int_{\mathbb{R}} (g'_* f' - g_* f) dv_* \right) d\theta,$$

where

$$\beta(\theta) \approx |\theta|^{-1-2s}, \quad 0 < s < 1.$$

The Kac equation for the fluctuation around the Maxwellian distribution reads

$$(3.1) \quad \begin{cases} \partial_t g + v \partial_x g + \mathcal{K}g = \Gamma(g, g), \\ g|_{t=0} = g_0, \end{cases}$$

*Theorem 3.2.* There exist some positive constants  $\epsilon_0 > 0$ ,  $c_0 > 1$  such that for all  $g_0 \in H^1(\mathbb{R}_x; L^2(\mathbb{R}_v))$  satisfying

$$\|g_0\|_{H^1(\mathbb{R}_x; L^2(\mathbb{R}_v))} \leq \epsilon_0,$$

the Cauchy problem (3.1) admits a unique weak solution  $g \in L^\infty([0, T]; H^1(\mathbb{R}_x; L^2(\mathbb{R}_v)))$ . Furthermore, this solution is smooth for all positive time  $0 < t \leq T$ , and satisfies the Gelfand-Shilov-Gevrey type estimates:

$$\forall \delta > 0, \exists C > 1, \forall 0 < t \leq T, \forall k, \ell, p \in \mathbb{N},$$

$$\|v^k \partial_v^\ell \partial_x^p g(t)\|_{L^\infty(\mathbb{R}_{x,v}^2)} \leq \frac{C^{k+l+p+1}}{t^{\frac{2s+1}{2s}(k+l+p+3)+\delta}} (k!)^{\frac{2s+1}{2s}} (\ell!)^{\frac{2s+1}{2s}} (p!)^{\frac{2s+1}{2s}} \|g_0\|_{H^1(\mathbb{R}_x; L^2(\mathbb{R}_v))}.$$

This result establishes a Gelfand-Shilov smoothing effect in the velocity variable and a Gevrey smoothing effect in the position variable :

$$\forall t > 0, \quad g(t, x, \cdot) \in S_{1+\frac{1}{2s}}^{1+\frac{1}{2s}}(\mathbb{R}_v)$$

and

$$\forall t > 0, g(t, \cdot, v) \in G^{1+\frac{1}{2s}}(\mathbb{R}_x).$$

So that there are similar regularity properties as Kolomogrov equation with some loss.

### The homogeneous Boltzmann equation with measure-valued initial datum

Recently, Y. Morimoto work for the homogeneous Boltzmann equation with measure-valued initial datum.

For  $\alpha \in [0, 2]$ ,  $\mathcal{P}^\alpha(\mathbb{R}^3)$  denotes the probability density function  $f$  on  $\mathbb{R}^3$  such that

$$\int_{\mathbb{R}^3} |v|^\alpha f(v) dv < \infty,$$

and moreover when  $\alpha \geq 1$ , it requires that

$$\int_{\mathbb{R}^3} v_j f(v) dv = 0, \quad j = 1, 2, 3.$$

A characteristic function  $\varphi(t, \xi)$  is the Fourier transform of  $f(t, v) \in \mathcal{P}^0(\mathbb{R}^3)$ :

$$\varphi(t, \xi) = \hat{f}(t, \xi) = \mathcal{F}(f)(t, \xi) = \int_{\mathbb{R}^3} e^{-iv \cdot \xi} f(t, v) dv.$$

For each  $\alpha \in [0, 2]$ , set  $\tilde{\mathcal{P}}^\alpha(\mathbb{R}^3) = \mathcal{F}^{-1}(\mathcal{K}^\alpha(\mathbb{R}^3))$  with  $\mathcal{K}(\mathbb{R}^3) = \mathcal{F}(\mathcal{P}^0(\mathbb{R}^3))$  and

$$\mathcal{K}^\alpha(\mathbb{R}^3) = \left\{ \varphi \in \mathcal{K}(\mathbb{R}^3) : \|\varphi - 1\|_{\mathcal{D}^\alpha} < \infty \right\}.$$

Here the distance  $\mathcal{D}^\alpha$  between  $\varphi(\xi)$  and  $\tilde{\varphi}(\xi)$  with  $\alpha > 0$  is defined by

$$\|\varphi - \tilde{\varphi}\|_{\mathcal{D}^\alpha} \equiv \sup_{0 \neq \xi \in \mathbb{R}^3} \frac{|\varphi(\xi) - \tilde{\varphi}(\xi)|}{|\xi|^\alpha}.$$

It is easy to see that

- The set  $\mathcal{K}^\alpha(\mathbb{R}^3)$  endowed with the distance  $\mathcal{D}^\alpha$  is a complete metric space;
- $\mathcal{K}^\alpha(\mathbb{R}^3) = \{1\}$  for all  $\alpha > 2$ ;
- the embeddings  $\{1\} \subset \mathcal{K}^\alpha(\mathbb{R}^3) \subset \mathcal{K}^\beta(\mathbb{R}^3) \subset \mathcal{K}(\mathbb{R}^3)$  hold for  $2 \geq \alpha \geq \beta \geq 0$ .

By taking the Fourier transform of the Boltzmann equation (Maxwellian case), we have the Bobylev formula:

$$(3.2) \quad \partial_t \varphi(t, \xi) = \int_{\mathbb{S}^2} \mathcal{B} \left( \frac{\xi \cdot \sigma}{|\xi|} \right) (\varphi(t, \xi^+) \varphi(t, \xi^-) - \varphi(t, \xi)) d\sigma,$$

where we have used

$$\varphi(t, 0) = \int_{\mathbb{R}^3} f(t, v) dv = 1,$$

here,

$$\xi^+ = \frac{\xi + |\xi|\sigma}{2}, \quad \xi^- = \frac{\xi - |\xi|\sigma}{2}$$

satisfying

$$\xi^+ + \xi^- = \xi, \quad |\xi^+|^2 + |\xi^-|^2 = |\xi|^2.$$

From now on, we consider the Cauchy problem for (3.2) with initial condition

$$(3.3) \quad \varphi(0, \xi) = \varphi_0(\xi).$$

Results available for the Cauchy problem (3.2)-(3.3):

- Global solvability: For  $\alpha \in ]2s, 2]$ , this Cauchy problem admits a unique global solution  $\varphi(t, \xi) \in C([0, \infty), \mathcal{K}^\alpha(\mathbb{R}^3))$  for every  $\varphi_0(\xi) \in \mathcal{K}^\alpha(\mathbb{R}^3)$
- Regularity: If  $\mathcal{F}^{-1}(\varphi_0)(v)$  is not a single Dirac mass,  $f(t, \cdot) \in L^1(\mathbb{R}^3) \cap H^\infty(\mathbb{R}^3)$  for any  $t > 0$  (see [5]).

When  $2s < \alpha < 2$ , the initial energy is infinite so that the solution will no longer tend to an equilibrium, but to a self-similar solution: For any given constant  $K > 0$

$$f_{\alpha, K}(t, v) = e^{-3\mu_\alpha t} \Psi_{\alpha, K}(ve^{-\mu_\alpha t}).$$

$$\mu_\alpha = \frac{\lambda_\alpha}{\alpha}, \quad \lambda_\alpha \equiv \int_{\mathbb{S}^2} \mathcal{B} \left( \frac{\xi \cdot \sigma}{|\xi|} \right) \left( \frac{|\xi^-|^\alpha + |\xi^+|^\alpha}{|\xi|^\alpha} - 1 \right) d\sigma,$$

$$\int_{\mathbb{R}^3} \Psi_{\alpha, K}(v) dv = 1, \quad \hat{\Psi}_{\alpha, K}(\xi) \in \mathcal{K}^\alpha(\mathbb{R}^3), \quad \lim_{|\eta| \rightarrow 0} \frac{1 - \hat{\Psi}_{\alpha, K}(\eta)}{|\eta|^\alpha} = K.$$

- Convergence to  $f_{\alpha,K}(t, v)$ : not well understood, even the convergence in distribution sense is a problem.

For  $f_0(v), g_0(v) \in \tilde{\mathcal{P}}^\alpha(\mathbb{R}^3)$ , set

$$\begin{aligned}\tilde{P}(t, \xi) &= e^{-At} \frac{1}{2} \sum_{j,l=1}^3 \xi_j \xi_l P_{jl}(0) X(\xi), \\ P_{jl}(0) &= \int_{\mathbb{R}^3} \left( v_j v_l - \frac{\delta_{jl}}{3} |v|^2 \right) (f_0(v) - g_0(v)) dv,\end{aligned}$$

where

$$A = \frac{3}{4} \int_{\mathbb{S}^2} \mathcal{B} \left( \frac{\sigma \cdot \xi}{|\xi|} \right) \left( 1 - \left( \frac{\sigma \cdot \xi}{|\xi|} \right)^2 \right) d\sigma,$$

$X(\xi) \equiv X(|\xi|)$  is a smooth radially symmetric function satisfying  $0 \leq X(\xi) \leq 1$  and  $X(\xi) = 1$  for  $|\xi| \leq 1$  and  $X(\xi) = 0$  for  $|\xi| \geq 2$ .

The first result on the  $\mathcal{D}^{2+\delta}$ -time asymptotic stability of the solutions is the following theorem.

*Theorem 3.3* (Morimoto-Yang-Zhao, *to appear JEMS*). Suppose  $f_0(v), g_0(v) \in \tilde{\mathcal{P}}^\alpha(\mathbb{R}^3)$  for  $\alpha \in ]\max\{2s, 1\}, 2]$ . Let  $\hat{f}(t, \xi)$  and  $\hat{g}(t, \xi)$  be the corresponding two global solutions of the Cauchy problem (3.2) with initial data  $\hat{f}_0(\xi)$  and  $\hat{g}_0(\xi)$  respectively. Assume for some  $\delta \in ]0, \alpha] \cap ]0, \frac{A}{\mu_\alpha}]$ , the initial data satisfy

$$(3.4) \quad \int_{\mathbb{R}^3} |v|^2 (f_0(v) - g_0(v)) dv = 0,$$

$$(3.5) \quad \begin{cases} \int_{\mathbb{R}^3} |v|^2 |f_0(v) - g_0(v)| dv < +\infty, \\ \left\| \hat{f}_0(\cdot) - \hat{g}_0(\cdot) - \tilde{P}(0, \cdot) \right\|_{\mathcal{D}^{2+\delta}} < +\infty. \end{cases}$$

Then there exists some positive constant  $C_1 > 0$  independent of  $t$  and  $\xi$  such that

$$\left\| \hat{f}(t, \cdot) - \hat{g}(t, \cdot) - \tilde{P}(t, \cdot) \right\|_{\mathcal{D}^{2+\delta}} \leq C_1 e^{-\eta_0 t}, \quad t \geq 0.$$

Here,  $\eta_0 = \min \{A - \delta \mu_\alpha, B\}$  and

$$B = \int_{\mathbb{S}^2} \mathcal{B} \left( \frac{\sigma \cdot \xi}{|\xi|} \right) \left( 1 - \left| \cos \frac{\theta}{2} \right|^{2+\delta} - \left| \sin \frac{\theta}{2} \right|^{2+\delta} \right) d\sigma, \quad \cos \theta = \frac{\sigma \cdot \xi}{|\xi|}.$$

Note that for the  $\mathcal{D}^{2+\delta}$ -convergence to the self-similar solution, one can simply take  $g_0 = \Psi_{\alpha,K}(v)$ . Based on this, in order to obtain a convergence in strong topology, such as in the Sobolev norms, we will give a uniform in time estimate on the solution in  $H^N$ -norm that is given in

*Theorem 3.4* (Morimoto-Yang-Zhao, *to appear JEMS*). For  $\max\{1, 2s\} < \alpha < 2$ , assume that  $f_0(v) \in \tilde{\mathcal{P}}^\alpha(\mathbb{R}^3)$  satisfies (3.4)-(3.5) and is not a single Dirac mass,  $g_0(v) = \Psi_{\alpha,K}(v)$ . Then for any given positive constant  $t_1 > 0$  and any  $N \in \mathbb{N}$ , there exists a positive constant  $C_2(t_1, N)$  such that

$$\sup_{t \in [t_1, +\infty)} \left\{ \|f(t, \cdot)\|_{H^N} \right\} \leq C_2(t_1, N).$$

Consequently, there exists a positive constant  $C_3(t_1, N)$  such that

$$(3.6) \quad \left\| f(t, \cdot) - f_{\alpha,K}(t, \cdot) \right\|_{H^N} = \left\| f(t, v) - e^{-3\mu_\alpha t} \Psi_{\alpha,K}(ve^{-\mu_\alpha t}) \right\|_{H^N} \leq C_3(t_1, N) e^{-\frac{\eta_0 t}{2}}$$

holds for any  $t \geq t_1$ .

the convergence rate given in (3.6) is faster than the decay rate of the self-similar solution itself. Hence in this case, the infinite energy solution  $f(t, v)$  converges to the self-similar solution  $f_{\alpha,K}(t, v)$  exponentially in time.

## References

- [1] Convergence to the self-similar solutions to the homogeneous Boltzmann equation, *to appear in J. Eur. Math. Soc.* (with Tong Yang and Huijiang Zhao)
- [2] Moment classification of infinite energy solutions to the homogeneous Boltzmann equation, *Analysis and Applications* (2015), on line doi:10.1142/S0219530515500232 663-683, (with Shuaikun Wang and Tong Yang)
- [3] Local existence of polynomial decay solutions to the Boltzmann equation for soft potentials, *Analysis and Applications* **13**(2015), 663-683, (with Tong Yang)
- [4] A new characterization and global regularity of infinite energy solutions to the homogeneous Boltzmann equation, *J. Math. Pures Appl.* **103** (2015), 809–829. (with S. Wang and T. Yang)
- [5] Smoothing effect of the homogeneous Boltzmann equation with measure-valued initial datum, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **32**(2015), 429-442, (with Tong Yang)
- [6] Gelfand-Shilov and Gevrey smoothing effect for the spatially inhomogeneous non-cutoff Kac equation. *J. Funct. Anal.* **269** (2015), 459-535. (with Nicolas Lerner, Karel Pravda-Starov and Chao-Jiang Xu)
- [7] Gelfand-Shilov smoothing properties of the radially symmetric spatially homogeneous Boltzmann equation without angular cutoff, *J. Differential Equations* **256** (2014), 797-831, (with Nicolas Lerner, Karel Pravda-Starov and Chao-Jiang Xu)
- [8] Local existence with mild regularity for the Boltzmann equation, *Kinet. Relat. Models* **6** (2013), 1011-1041, (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)

- [9] A remark on the ultra-analytic smoothing properties of the spatially homogeneous Landau equation, *Kinet. Relat. Models* **6** (2013), 715-727, (with Karel Pravda-Starov and Chao-Jiang Xu)
- [10] Spectral and phase space analysis of the linearized non-cutoff Kac collision operator, *J. Math. Pures Appl.*(9)**100** (2013), 832-867, (with Nicolas Lerner, Karel Pravda-Starov and Chao-Jiang Xu)
- [11] Phase space analysis and functional calculus for the linearized Landau and Boltzmann operators, *Kinet. Relat. Models***6** (2013), 625-648, (with Nicolas Lerner, Karel Pravda-Starov and Chao-Jiang Xu)
- [12] A remark on Cannone-Karch solutions to the homogeneous Boltzmann equation for Maxwellian molecules, *Kinet. Relat. Models*, **5** (2012), 551-561 .
- [13] Smoothing effect of weak solutions for the spatially homogeneous Boltzmann Equation without angular cutoff, *Kyoto J. Math.*,**52** (2012) , 433-463, (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
- [14] Hypoelliptic estimates for a linear model of the Boltzmann equation without angular cutoff, *Comm. Part. Diff. Equat.*, **37** (2012), 234-284, (with Nicolas Lerner and Karel Pravda-Starov)
- [15] The Boltzmann equation without angular cutoff in the whole space: I, Global existence for soft potential, *J. Funct. Anal.*, **262** (2012), 915-1010, (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
- [16] Uniqueness of solutions for the non-cutoff Boltzmann Equation with soft potential, *Kinet. Relat. Models*, **4-4** (2011), 919-934. (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
- [17] The Boltzmann equation without angular cutoff in the whole space: Qualitative properties of solutions, *Arch. Rational Mech. Anal.*, 202(2011),599-661, (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
- [18] The Boltzmann equation without angular cutoff in the whole space: II, global existence for hard potential, *Analysis and Applications*, **9-2** (2011), 113-134. (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
- [19] Bounded solutions of the Boltzmann equation in the whole space, *Kinetic and Related Models*,**4** (2011) 17-40, (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
- [20] Global existence and full regularity of the Boltzmann equation without angular cutoff, *Comm. Math. Phys.* ,**304-2**(2011),513-581 (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
- [21] Regularizing effect and local existence for non-cutoff Boltzmann equation, *Arch. Rational Mech. Anal.*,**198** (2010), 39-123, (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
- [22] Gevrey smoothing effect of solutions for spatially homogeneous nonlinear Boltzmann equation without angular cutoff, *J. Pseudo-Differ. Oper. Appl.***1** (2010), 139-159, (with Seiji Ukai)
- [23] Instability of the Cauchy-Kovalevskaya solution for a class of non-linear systems, *American Journal Mathematics*,**132(1)**(2010) 99-123, (with Nicolas Lerner and Chao-Jiang Xu)
- [24] Regularity of solutions for the Boltzmann equation without angular cutoff. *C. R. Math. Acad. Sci. Paris, Ser. I*, **347** (2009), no. 13-14, 747-752. (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
- [25] Local existence for non-cutoff Boltzmann equation, *C. R. Math. Acad. Sci. Paris, Ser. I*,



- 347** (2009), no. 21-22, 1237–1242. (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
- [26] Ultra-analytic effect of Cauchy problem for a class of kinetic equations, *J. Differential Equations*, **247**(2009) 596-617, (with Chao-Jiang Xu)
  - [27] Regularity of solutions to the spatially homogeneous Boltzmann equation without angular cutoff, *Discrete and Continuous Dynamical Systems - Series A*, **24**(2009), 187-212, (with Seiji Ukai, Chao-Jiang Xu and Tong Yang)
  - [28] Uncertainty principle and kinetic equations, *J. Funct. Anal.*, **255** (2008) 2013-2066, (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
  - [29] Regularity of solutions for spatially homogeneous Boltzmann equation without Angular cutoff. *Kinetic and Related Models*, **1** (2008) 453-489, (with Zahoui Huo, Seiji Ukai, Tong Yang)
  - [30] Uncertainty principle and regularity for Boltzmann type equations, *C.R.Acad. Sci. Paris, Ser. I*, **345**, 667-672 (2007) (with Radjesvarane Alexandre, Seiji Ukai, Chao-Jiang Xu and Tong Yang)
  - [31] Hypoellipticity for a class of kinetic equations, *J. Math. Kyoto Univ.* **47-1**, 129-152 (2007) (with Chao-Jiang Xu)
  - [32] On the Fefferman-Phong inequality and a Wiener-type algebra of pseudodifferential operators, *Publ. RIMS Kyoto Univ.* , **43**, 329-371 (2007) (with Nicolas Lerner).
  - [33] Nonlinear hypoellipticity of infinite type, *Funkciaj Ekvacioj*, **50**, 33-65 (2007) (with Chao-Jiang Xu).
  - [34] The collision term of Boltzmann Equation for Debye-Yukawa potential, *Seminar Notes of Mathematical Sciences, Ibaraki Univ.*, **9**. 90-99 (2006) (with Seiji Ukai, Chao-Jiang Xu and Tong Yang)
  - [35] A Wiener algebra for the Fefferman-Phong inequality *Seminaire Équations Aux Dérivées Partielles* **2005-2006**, 12 pp., *Ecole Polytechnique*, (with Nicolas Lerner).
  - [36] A viewpoint on positivity of pseudodifferential operators from the Wick calculus *Recent Trends in Microlocal Analysis, RIMS Kokyuroku* **1412**, 80–90 (2005) (with Nicolas Lerner)
  - [37] Regularity of weak solutions for a class of infinitely degenerate elliptic semilinear equations, *Seminaire Équations Aux Dérivées Partielles* **2003-2004**, 12 pp., *Ecole Polytechnique*, (with Chao-Jiang Xu).
  - [38] Logarithmic Sobolev inequality and semi-linear Dirichlet problems for infinitely degenerate elliptic operators, *Astérisque*, **284**, 245–264 (2003). (with Chao-Jiang Xu).
  - [39] Wick calculus and the Cauchy problem for some dispersive equations, *Osaka J. Math.*, **39-1**, 123–147 (2002) ( with Hiroshi Ando).
  - [40] Remark on the analytic smoothing for the Schrödinger equation, *Indiana Univ. Math. J.* **49-4**, 1563-1579(2000) ( with Luc Robbiano and Claude Zuily).
  - [41] Hypoellipticity for elliptic operators with infinite degeneracy, "Partial Differential Equations and Their Applications" (Chen Hua and L. Rodino, eds.), *World Sci. Publishing, River Edge, NJ*, 240-259(1999) ( with Tatsushi Morioka).
  - [42] Hypoellipticity for operators of infinitely degenerate Egorov type, *Tsukuba J. Math.* **23-1**, 215-224(1999). (with Nils Dencker and Tatsushi Morioka).
  - [43] Hypoellipticity for a class of infinitely degenerate elliptic operators "Geometrical Optics and Related Topics"(F. Colombini and N. Lerner, eds.), *Birkhäuser, Boston, Mass., USA*, 295-317(1997) ( with Tatsushi Morioka).
  - [44] The positivity of Schrödinger operators and the hypoellipticity of second order degenerate elliptic operators, *Bull. Sc. Math.* **121**, 507-547(1997) ( with Tatsushi Morioka).

- [45] Local solvability and hypoellipticity for pseudodifferential operators of Egorov type with infinite degeneracy, *Nagoya Math. J.* **139**, 151-171(1995).
- [46] Hypoelliptic operators of principal type with infinite degeneracy, *Tsukuba J. Math.* **19-1**, 187-200(1995).
- [47] Some remarks on hypoelliptic operators which are not micro-hypoelliptic *Publ. RIMS Kyoto Univ.* **28** 579-586(1992) ( with Tatsushi Morioka).
- [48] Hypoelliptic operators in  $R^3$  of the form  $X_1^2 + X_2^2$ , *J. Math. Kyoto Univ.* **32** 461-484(1992).
- [49] Estimates for degenerate Schrödinger operators and hypoellipticity for infinitely degenerate elliptic operators, *J. Math. Kyoto Univ.* **32** 333-372(1992) .
- [50] Estimates for degenerate Schrödinger operators and an application for infinitely degenerate hypoelliptic operators, *Proc. Japan Acad.*, **65**, 155-157 (1989).
- [51] Propagation of wave front set and hypoellipticity for degenerate elliptic operators, *Pitman Res. Notes Math. Ser.* **183**, 212-224 (1988).
- [52] The uncertainty principle and hypoelliptic operators, *Publ. RIMS Kyoto Univ.* **23** , 955-964 (1987).
- [53] Criteria for hypoellipticity, *Seminaire Équations Aux Dérivées Partielles 1986-1987*, *Ecole Polytechnique*.
- [54] A criterion for hypoellipticity of second order differential operators, *Osaka J. Math.*, **24** , 651-675 (1987).
- [55] Hypoellipticity for infinitely degenerate elliptic operators, *Osaka J. Math.*, **24** , 13-35 (1987).
- [56] Criteria for hypoellipticity of differential operators, *Publ. RIMS Kyoto Univ.* **22** , 1129-1154 (1986).
- [57] Propagation of wave front sets of solutions of the Cauchy problem for a hyperbolic system in Gevrey classes, *Osaka J. Math.*, **23** , 765-814 (1986) ( with Kazuo Taniguchi ).
- [58] Non-hypoellipticity for degenerate elliptic operators, *Publ. RIMS Kyoto Univ.* **22** , 25-30 (1986).
- [59] On a criterion for hypoellipticity, *Proc. Japan Acad.*, **62**, 137-140 (1986).
- [60] Propagation of wave front set in Gevrey class for an example of hyperbolic system, *Nagoya Math. J.* **101**, 131-150 (1986).
- [61] Precise propagation of singularities for a hyperbolic system with characteristics of variable multiplicity, *Nagoya Math. J.* **101**, 111-130 (1986) ( with Chisato Iwasaki ).
- [62] Propagation of wave front sets of solutions of the Cauchy problem for a hyperbolic system in Gevrey classes, *Proc. Japan Acad.*, **61**, 27-30 (1985) ( with Kazuo Taniguchi ).
- [63] Propagation of singularities for a hyperbolic system with nilpotent characteristics, II, *Comm. Partial Differential Equations* **9**, 1407-1436 (1984) ( with Chisato Iwasaki ).
- [64] Opérateur pseudodifférentiel et classes de Gevrey, *Comm. Partial Differential Equations* **8**, 237-255 (1983) ( with Syuji Hashimoto and Tadato Matsuzawa ).
- [65] On hypoelliptic operators with multiple characteristics of odd order, *Osaka J. Math.*, **20**, 237-255 (1983).
- [66] Propagation of singularities for a hyperbolic system with nilpotent characteristics, I, *Comm. Partial Differential Equations* **7**, 743-794 (1982) ( with Chisato Iwasaki ).
- [67] On the propagation of the wave front set of a solution for a hyperbolic system, *Math. Japonica* **27**, 501-508 (1982).
- [68] Fundamental solution for a hyperbolic equation with involutive characteristics of variable multiplicity, *Comm. Partial Differential Equations* **4**, 609-643 (1979).

- [69] On the hypellipticity for infinitely degenerate semi-elliptic operators, *J. Math. Soc. Japan* **30**, 327-358 (1978).